

PERIODIC MONITORING OF MARINE ENVIRONMENT FOR MSPL OUTFALL, BHAVNAGAR, GUJARAT

**Monitoring Report - 3 of 4/2021-22
November 2021**

PROJECT CODE: 656041920



**MADHU SILICA PRIVATE LIMITED
BHAVNAGAR, GUJARAT**



INDOMER COASTAL HYDRAULICS (P) LTD.
(ISO 9001: 2015 CERTIFIED, QCI - NABET & NABL)
63, GANDHI ROAD, ALWAR THIRUNAGAR, CHENNAI 600 087.
Tel: + 91 44 2486 2482 to 84; M: + 91 99401 41650; Fax: + 91 44 2486 2484
Web site: www.indomer.com, E-mail: ocean@indomer.com



INDOMER COASTAL HYDRAULICS (P) LTD.

(ISO 9001 : 2015 CERTIFIED,QCI-NABET, NABL & CDC-MoST ACCREDITED)

63, Gandhi Road, Alwar Thirunagar, Chennai 600 087.

Tel: + 91 44 2486 2482 to 84; M: + 91 99401 41650; Fax: + 91 44 2486 2484

Web site: www.indomer.com, E-mail: ocean@indomer.com

Client	Madhu Silica Private Limited, Bhavnagar.				
Project Title	Periodic Monitoring of Marine Environment for MSPL Outfall, Bhavnagar, Gujarat for the year 2021-22.				
Project Code	656041920				
Abstract	<p>Madhu Silica Private Limited (MSPL) is discharging 10 MLD treated industrial effluent in Bhavnagar Creek for which MoEFCC-CRZ clearance was obtained vide F. No. 11-6/2015-IA-III dt. 29.06.2015.</p> <p>Subsequently, MSPL has been submitting the Periodic Monitoring Reports every 3 months.</p> <p>Status of reports for year 2021-22:</p> <ul style="list-style-type: none"> May 2021 – 1 of 4 – Report not done due to COVID 19 Pandemic second wave. Sampling could not be done due to transport restrictions. Aug 2021 – 2 of 4 – Report Submitted 12.08.2021 Nov 2021 – 3 of 4 – Present report. <p><i>*Due to COVID-19 Nationwide lockdown and strict interstate travel protocols, quarterly monitoring study for the 1st season period March 2021 – May 2021 was not conducted. However, after the interstate relaxations, 2nd season quarterly monitoring study for June 2021 to August 2021 was conducted and report has been submitted. This forms as the 3rd season Quarterly Monitoring Report (September to November 2021) for the year 2021-22.</i></p>				
Document type	Controlled				
References	W.O. Email dt. 23/04/2019				
Date	Report Type	Originator	Checked by	Approved by	Approver's sign
18.11.2021	Final	Mr. J. Jaganmohan	Mr. J. Guru Prasath	Dr. P. Chandramohan	
1	Project Code	656041920		Text pages	25
2	File Location	\\192.168.1.200/Indomer/MSPL-Nov.21/Report		Tables	19
				Figures	2

CONTENT

	Page
Content	i
List of Tables	ii
List of Figures	iii
1 PROJECT DESCRIPTION	1
2 SCOPE OF MONITORING	2
3 DATA COLLECTION FOR MONITORING	3
4 RESULTS	8
4.1 Water quality	8
4.2 Sediment quality	9
4.3 Biological parameters	9
5 ENVIRONMENTAL STATUS - COMPARISON WITH PRE-PROJECT PERIOD	11
Annexure I – QCI NABET accreditation certificate	
Annexure II – NABL accreditation certificate	

LIST OF TABLES

Table

- 1 Details of sampling locations
- 2 Water quality parameters at Bhavnagar creek (November 2021)
- 3 Biochemical Oxygen Demand and Chemical Oxygen Demand at Bhavnagar creek water (November 2021)
- 4 Concentration of Heavy Metals, Phenols and Total Petroleum Hydrocarbons at Bhavnagar creek water (November 2021)
- 5 Sediment texture at Bhavnagar creek (November 2021)
- 6 Sediment quality parameters at Bhavnagar creek (November 2021)
- 7 Concentration of Heavy Metals, Phenols and Total Petroleum Hydrocarbons in sediments at Bhavnagar creek (November 2021)
- 8 Primary productivity at Bhavnagar creek water (November 2021)
- 9 Phytoplankton species composition at Bhavnagar creek water (November 2021)
- 10 Phytoplankton numerical abundance (cells/l) at Bhavnagar creek water (November 2021)
- 11 Phytoplankton population at Bhavnagar creek water (November 2021)
- 12 Numerical abundance of zooplankton (nos./100m³) at Bhavnagar creek water (November 2021)
- 13 Zooplankton diversity abundance and biomass at Bhavnagar creek water (November 2021)
- 14 Subtidal and Intertidal benthic population at Bhavnagar creek (November 2021)
- 15 Bacterial population at Bhavnagar creek water ($\times 10^3$ CFU/ml) (November 2021)
- 16 Bacterial population at Bhavnagar creek sediments ($\times 10^4$ CFU/g) (November 2021)
- 17 Comparison of Water quality parameters between Pre-project period (May 2013) and Monitoring period (November 2021)
- 18 Comparison of Sediment quality parameters between Pre-project period (May 2013) and Monitoring period (November 2021)
- 19 Comparison of Biological parameters between Pre-project period (May 2013) and Monitoring period (November 2021)



LIST OF FIGURES

Figure

- 1 Location map
- 2 Sampling locations

1. PROJECT DESCRIPTION

Madhu Silica Pvt. Ltd. (MSPL), Bhavnagar, is the largest manufacturer of precipitated Silica in India and it is the 3rd largest company in the world. MSPL with its allied company Aqua gel Chemicals Pvt. Ltd., is having seven established plants capable of manufacturing around 95000 MT of precipitated Silica per annum. It has recently established a state of art plant with capacity of 45000 MT/Annum in Bhavnagar. The company has a large business associate network in India and Asia, Middle East, Latin America, US & Europe. Apart from the large Indian market, the company exports their products to more than 40 countries. The company has on its credit ISO 9001:2008, ISO 14001:2004, OHSAS 18001:2007, FAMI-QS, FDA certified and registered for each.

MSPL has planned for capacity expansion in 28 acres of land towards total installed capacity of precipitate Silica of 95000 MT/Annum. Under this development process, MSPL has obtained permission to discharge additional 10 MLD (417 m³/hour) treated industrial effluent in the marine environment. It has laid a submarine pipeline with diffuser on the creek bed for discharging the effluent.

MSPL obtained MoEFCC clearance vide F. No. 11-6/2015-IA-III dt. 29.06.2015. The post project monitoring is being carried out for four seasons in a year to comply the conditions of MoEFCC covering the aspects of seawater quality, seabed sediment quality and marine benthic flora & fauna. The monitoring studies are being carried out from May 2017 i.e. since the commencement of project on quarterly basis. The periodic monitoring reports are being regularly submitted to the Regional Office of MoEFCC, Bhopal. This report forms as the third periodic monitoring (November 2021) for the year 2021 - 2022.

[Due to COVID-19 Nationwide lockdown and strict interstate travel protocols, quarterly monitoring study for the 1st season period March 2021 – May 2021 was not conducted. However, after the interstate relaxations, 2nd season quarterly monitoring study for June 2021 to August 2021 was conducted and report has been submitted. This forms as the 3rd season Quarterly Monitoring Report \(September to November 2021\) for the year 2021-22.](#)

The location map is shown in Fig. 1 and the effluent outfall location is shown in Fig. 2.

2. SCOPE OF MONITORING

Periodic monitoring near the outfall location as suggested in the Post Project Monitoring Plan of the EIA report was recommended by the MoEFCC. Accordingly, MSPL has asked Indomer Coastal Hydraulics (P) Ltd, Chennai, to take up periodic monitoring programme. It was proposed to conduct quarterly monitoring covering four times in a year i.e., in the months of May, August, November and February every year during the project operational period.

Indomer Coastal Hydraulics (P) Ltd., Chennai is an ISO 9001:2015 organization, NABL and QCI - NABET accredited organization vide NABET/EIA/2023/RA 0207 dt. 29.06.2021 for Sector 27: Oil & Gas Transportation pipeline (crude and refinery/petrochemical products), passing through national parks/ sanctuaries/ coral reefs/ ecologically sensitive areas including LNG Terminal and Sector 33: Ports, harbours, jetties, marine terminals, breakwaters and dredging.

QCI-NABET accreditation certificate is attached as Annexure I.

3. DATA COLLECTION FOR MONITORING

The creek water, creek bed sediments and biological samples were collected at three locations around the outfall points (SS1 to SS3) in November 2021. The samples for intertidal benthos were also collected at three locations (IB1 to IB3). The details of the sampling locations are given in Table 1 and shown in Fig. 2. The monitoring work covered is listed below:

i) Creek water quality parameters at three locations at surface and bottom:

Temperature
pH
Salinity
Dissolved Oxygen (DO)
Total Dissolved Solids (TDS)
Biochemical Oxygen Demand (BOD)
Chemical Oxygen Demand (COD)
Ammonium
Nitrite
Nitrate
Total nitrogen
Phosphate
Total phosphorus
Turbidity
Total Suspended Solids (TSS)
Cadmium
Lead
Mercury
Total Chromium
Phenols and
Total Petroleum Hydrocarbons

ii) Creek bed sediment quality parameters at three locations:

Sediment Texture
Total Nitrogen
Total Phosphorous
Total organic carbon
Calcium carbonate
Cadmium
Lead
Mercury
Total Chromium
Phenols and
Total Petroleum Hydrocarbons

iii) Biological parameters at three locations on:

Primary Productivity

Phytoplankton, its biomass and diversity

Zooplankton, its biomass and diversity

Sub-tidal and inter-tidal macrobenthos, its biomass and diversity and

Microbial population in creek water and sediment

Analysis: All the water and sediment parameters were analyzed by Indomer Coastal Hydraulics Pvt. Ltd., which is accredited by the National Accreditation Board for Testing and Calibration Laboratories (NABL) vide certificate no. TC-5232/12.01.2022. Analysis of Total Petroleum Hydrocarbons in creek water and sediment samples was outsourced to NABL accredited Laboratory. NABL accreditation certificate of Indomer is attached as Annexure II.

Sample Analysis Protocol: Creek water and creek bed sediment samples were analyzed as per the IS/APHA/USEPA standard methods, details of which are given below.

Sl.No.	Parameters	Protocol
Water quality		
1	Temperature	IS 3025 : Part 9
2	pH	IS 2720 : Part 11
3	Salinity	IND/SOP/WQ/13
4	Dissolved Oxygen	IS 3025 : Part 38
5	Total Dissolved Solids	IS 3025 : Part 16
6	BOD	IS 3025 : Part 44
7	COD	IS 3025 : Part 58
8	Turbidity	IS 3025 : Part 10
9	Ammonium	IS 3025 : Part 34
10	Nitrite	IS 3025 : Part 34
11	Nitrate	IS 3025 : Part 34
12	Phosphate	IS 3025 : Part 31
13	Total Nitrogen	IS 3025 : Part 34
14	Total Phosphorous	IS 3025 : Part 31
15	Total Suspended Solids	IS 3025 : Part 17
16	Cadmium	IS 3025 : Part 2
17	Lead	IS 3025 : Part 2
18	Chromium	IS 3025 : Part 2
19	Mercury	IND/SOP/WQ/35
20	Phenols	IS 3025 : Part 43
21	Petroleum Hydrocarbons	TNRCC method 1055
Sediment quality		
1	Soil texture	IS 2720 : Part 4
2	Total Organic Carbon	IS 2720 : Part 22
3	Total Nitrogen	IS 14684 – 1999
4	Total Phosphorous	IS 10158 – 1982
5	Calcium carbonate	IS 2720 : Part 23
6	Cadmium	USEPA 3050 B
7	Lead	USEPA 3050 B
8	Chromium	USEPA 3050 B
9	Mercury	USEPA 3050 B
10	Phenols	USEPA 8041 & 3545 A
11	Petroleum Hydrocarbons	TNRCC method 1055

Biological parameters

Primary Productivity: Primary production will be estimated from appropriate study area sampling stations. From the water sampler, the samples must be immediately transferred to 125 ml DO bottles (two light bottles and one dark bottle). The sample in the first bottle will be used immediately to determine the initial level of dissolved oxygen (DO) content followed by Winkler method. The light and dark bottles will be incubated under water for a period of 6 hr and dissolved oxygen will be measured. Primary productivity was calculated by oxygen method. Oxygen values will be converted to carbon values by applying the equation.

Phytoplankton: Phytoplankton samples will be collected from appropriate study area sampling stations, for both qualitative and quantitative analyses.

Phytoplankton samples for quantitative analyses will be taken by 1 liter plastic container from surface water and preserved with Lugol's iodine solution. The analysis of phytoplankton samples include initial concentration of water sample to 15 ml volume based on settling and siphoning procedure. Quantitative estimation of phytoplankton will be done by counting in Sedge wick-Rafter cell counter. It involved calculation of the number of cells of each species of phytoplankton in one liter of water sample.



For the qualitative analysis, phytoplankton samples will be collected using circular standard plankton net (60 μ mesh and 60 cm mouth diameter). The net will be towed at subsurface for 5 minutes. After the collection, samples must be preserved in 4% buffered formaldehyde and analyzed under an inverted microscope following the standard literature (R. Subrahmanyam, 1946; C.P. Gopinathan, 1976 and Thomas, 1997).

Zooplankton: Zooplankton samples will be collected using circular zooplankton net (300 μ mesh and 60 cm mouth diameter). The samples must be collected during day time to calculate their biomass, population and bio diversity. The net will be towed for 5 minutes. After the collection, samples must be preserved in 5% buffered formaldehyde. The biomass value of zooplankton will be calculated using the displacement volume method. The faunal composition and the relative abundance of different zooplankton taxa will be sorted out and identified from aliquots upto species level as far as possible. All taxonomic observation and measurements will be made on preserved samples. Specimens will be identified based on the standard manuals (Kasturirangan, 1963; and Conway et al. 2003). The estimated abundance (density) for the different groups will be expressed as nos. /100m³.

Flowmeter: Digital Flowmeter (model - 2030R) duly calibrated by the company will be used for estimating the volume of flow into the net towed for 5 minutes for the collection of phytoplankton and zooplankton. The flow meter consists of an impeller and a counter. The impeller is directly connected to the counter which records each revolution of the impeller. The flow meter has to be attached to the mouth region of the plankton net.

Macro Benthos: Seabed sediment samples will be collected using Van Veen grab from sampling stations. The intertidal benthic samples will be collected from appropriate stations. The benthic organisms will be separated by sieving through 500 micron mesh and preserved using formaldehyde and Rose Bengal stain. The samples will be sorted and identified upto

groups/genera level using stereo microscope. The wet weight will be taken to calculate the biomass of benthic organisms.

Microbiology: The microbiological samples will be collected from appropriate study area sampling stations. The total coliform from each location will be identified by membrane filter technique (APHA 9060 A & B). Samples will be collected clean, sterile and non-reactive glass or plastic bottles. Microbial analysis is started as soon as possible after collection to avoid unpredictable changes. Spread plate method will be used to culture the microorganisms. The agar media used for analysis were: Nutrient agar, MacConkey agar, Thiosulphate Citrate Bile Sucrose agar, Xylose Lysine Deoxycholate agar, M-Enterococcus agar and Cetrimide agar. Plates were incubated at 37° C for 48 hrs. After incubation, the colonies will be counted and identified based on their colour characteristics.

4. RESULTS

4.1. Water quality

The estimated creek water quality parameters on temperature, pH, salinity, total dissolved solids, dissolved oxygen, ammoniacal-nitrogen, nitrite-nitrogen, nitrate-nitrogen, total nitrogen, phosphate-phosphorus, total phosphorus, total suspended solids, and turbidity are presented in Table 2. Biochemical Oxygen Demand and Chemical Oxygen Demand are presented in Table 3. Results of the heavy metals cadmium, lead, total chromium, mercury, phenols and total petroleum hydrocarbons are presented in Table 4.

During the present study, creek water salinity ranged from 19.1 to 20.1 PSU and pH range was 7.34 to 7.49. The values of salinity and pH are very low compared to previous study, which may be due to freshwater influx and effect of monsoon rain runoff in the creek water. Very high amount of fresh water flow was observed in the creek during the sample collection. Hence, salinity and pH values were low in the creek water.

The range of TSS (1520 to 1786 mg/l) was found to be high compared to the previous sampling period of august 2021. Dissolved oxygen values ranged from 5.8 to 6.1 mg/l, which is a small fluctuation, to the earlier reported values. BOD values were also found to be normal (<2 mg/l) like earlier periods at all the three stations.

Ammoniacal-Nitrogen, Nitrite, Nitrate, Total Nitrogen, Phosphate and Total Phosphorus concentration levels show seasonal variation and however all the nutrients values are observed to be within the normal range. Cadmium levels ranged from 1.95 to 2.01 µg/l and Chromium concentration ranged from 2.08 to 2.26 µg/l. However, the metals were observed to be in trace level in the study areas. Similarly, phenols and total petroleum hydrocarbons were also found to be below detectable levels during the present study.

Creek water values for Turbidity, TSS, DO, BOD, nutrients, heavy metals and other organic compounds were found to be in normal range. Hence, it can be confirmed that treated effluent is discharged into the creek environment do not affect the creek water quality.

4.2. Sediment quality

The creek bed sediment quality parameters collected at 3 locations (SB1, SB2 and SB3) are given in Tables 5 and 6. Results of cadmium, lead, total chromium, mercury, phenols and total petroleum hydrocarbons in sediments are presented in Table 7. Sediment texture, fine sand nature was found during the present study period, at all stations. Total phosphorus, calcium carbonate, total nitrogen and total organic carbon values at all stations, varied slightly, compared to pre-project period study. Total nitrogen values ranged from 26.8 to 42.6 mg/kg at the three stations. Cadmium, mercury, lead, phenols and total petroleum hydrocarbon levels were found to be below detectable limit, at all the three stations. However, marginal difference was observed in the chromium level, between the pre-project period and the present study (23.2 to 24.6 mg/kg).

During present study, sediment nature was found to be fine sand texture. The levels of trace metals mercury and lead at all three stations along with phenols and petroleum hydrocarbons in the sediment were found to be below detectable level. Chromium values ranged from 23.2 to 24.6 mg/kg at all three stations. In general, these conditions reflect that the creek bed sediments are in normal condition and remain uncontaminated.

4.3. Biological parameters

Phytoplankton and primary productivity: Phytoplankton are the primary source of food in the marine environment. The concentration and numerical abundance of the phytoplankton indicate the fertility of a region. The phytoplankton population depends primarily upon the nutrients present in the creek water and the sunlight for photosynthesis. This primary production is an important source of food, for the higher organisms, in the marine environment. The measured primary production results indicate that the area is moderately productive, and the values varied between 330 and 420 mgC/m³/day; the recorded average value is 370 mgC/m³/day (Table 8).

Various phytoplankton groups were observed, and their percentage composition and numerical abundance are shown in Table of 9 and 10. Phytoplankton diversity fluctuated from 21 to 29 species. Bacillariophyceae (diatoms) formed the major group followed by Dinophyceae (dinoflagellates) and Cyanophyceae (blue green algae). Phytoplankton population density varied from 2400 to 2750 cells/l (Table 11).

Phytoplankton population were found three classes namely, Bacillariophyceae (75.6%), Dinophyceae (15.4%) and Cyanophyceae (9.0 %) (Table 10). During the study period, *Coscinodiscus centralis* was the most dominant species in the study area followed by *Odontella mobiliensis*, *Ceratium sp*, *Bacillaria paradoxa* and *Triceratium sp*.

Zooplankton: The numerical abundance of zooplankton varied from 8275 to 10594 nos./100m³ (Table 12). The highest zooplankton population was observed at SS2 and the lowest was observed at SS3. The zooplankton biomass varied from 10.7 to 13 ml/100m³ (Table 13). It is observed in seasonal variation. The most dominant zooplankton species are *Oithona nana*, *Brachionus plicatilis*, Copepod nauplii and *Nannocalanus minor* among the total of 29071 species.

Benthos: Benthic faunal population in an environment depends on the nature of the substratum and the organic matter content of the substratum.

Sub-tidal benthos: The numerical abundance of the benthic fauna varied from 180 to 520 nos./m² (Table 14).

Inter-tidal benthos: The existence of fauna appeared to be moderate in the three (IB1 to IB3) locations. The numerical abundance of the intertidal benthic fauna varied between 180 and 240 nos./m² and the intertidal faunal population is shown in Table 14.

Microbiology:

Bacterial counts in the water and sediment samples were analyzed and presented in Tables 15 and 16 respectively. In the water samples, population density ranged from 0.01 to 5.18×10³ CFU/ml. In the sediment, the population density ranged from 0.02 to 5.23×10⁴ CFU/g. Bacterial population was found normal. In general, there is not much of a variation observed between the present and earlier values both in creek water and sediment.

Primary production, phytoplankton and zooplankton are found to be in normal values. The sub-tidal and inter-tidal benthic population showed normal range at all three stations during the study. Bacterial population in water and sediments also indicated normal range.

5. ENVIRONMENTAL STATUS - COMPARISON WITH PRE-PROJECT PERIOD

Comparison of water quality, sediment quality and biological parameter results were carried out between the pre-project period (May 2013) and the periodic monitoring results of November 2021 as given in Tables 17, 18 and 19.

Water Quality

It is inferred that the water quality parameters showed normal range as applicable for the creek waters in Bhavnagar and they are comparable with the pre-project period results. The differences are very marginal, and they are due to seasonal variation which is common for creek waters.

Sediment quality

Sediment texture was predominantly towards fine sand in nature during present study and other chemical properties are observed to be in normal range and there is no significant change in the creek bed sediment quality.

Biological Parameters

Primary production, Phytoplankton and Zooplankton population were showing marginal seasonal variation but it notified within the normal range in the present observation. Also found the attribute range meets the optimum water quality conditions. The sub-tidal and intertidal benthic population showed normal range compare than previous monitoring period of November 2021.

Microbiology

There are no changes observed in bacterial population, in water and sediments, among the different sampling periods in this creek.

Table 1. Details of sampling locations

Station	UTM Coordinates (WGS 84)		Water depth (m)	Sampling depth* (m)
	X (m)	Y (m)		
WATER SAMPLING LOCATIONS				
SS1	204022	2414736	2.7	Surface & Bottom
SS2	204458	2414511	2.9	Surface & Bottom
SS3	204951	2414447	3.1	Surface & Bottom
SEABED SEDIMENT SAMPLING LOCATIONS				
SB1	204022	2414736	-	-
SB2	204458	2414511	-	-
SB3	204951	2414447	-	-
SUB-TIDAL BENTHOS SAMPLING LOCATIONS				
SB1	204022	2414736	-	-
SB2	204458	2414511	-	-
SB3	204951	2414447	-	-
INTERTIDAL BENTHOS SAMPLING LOCATIONS				
IB1	204178	2414637	Intertidal zone	
IB2	204435	2414473	Intertidal zone	
IB3	204650	2414381	Intertidal zone	

*SS =Seawater Sample, SB = Seabed Sediment

Table 2. Water quality parameters at Bhavnagar creek (November 2021)

Station	Water depth*	Temp. (°C)	Salinity (PSU)	pH	DO (mg/l)	TSS (mg/l)	Turbidity (NTU)	Ammoniacal Nitrogen (μmol/l)	Nitrite (μmol/l)	Nitrate (μmol/l)	Total Nitrogen (μmol/l)	Phosphate (μmol/l)	Total Phosphorus (μmol/l)
SS1	Surface	26.5	19.1	7.34	6.0	1628	740	2.27	1.20	5.7	9.26	2.3	3.4
	Bottom	26.0	19.4	7.48	5.9	1786	812	1.96	1.29	6.6	9.94	2.1	3.0
SS2	Surface	26.7	19.4	7.42	6.1	1598	726	2.05	1.28	4.2	7.62	2.2	3.4
	Bottom	26.2	19.7	7.47	5.9	1742	792	1.77	1.34	5.2	8.42	2.0	2.4
SS3	Surface	27.0	19.7	7.45	6.0	1520	691	1.94	1.26	4.3	7.55	1.6	3.0
	Bottom	26.5	20.1	7.49	5.8	1688	767	1.61	1.29	5.6	8.56	1.5	2.5

Table 3. Biochemical Oxygen Demand and Chemical Oxygen Demand at Bhavnagar creek water (November 2021)

Station	Water depth	BOD (mg/l)	COD (mg/l)
SS1	Surface	1.4	30.4
	Bottom	1.6	33.6
SS2	Surface	1.3	28.8
	Bottom	1.5	32.0
SS3	Surface	1.2	25.6
	Bottom	1.4	27.2

Table 4. Concentration of Heavy Metals, Phenols and Total Petroleum Hydrocarbons at Bhavnagar creek water (November 2021)

Station	Water depth	Heavy metals (µg/l)				Phenols (mg/l)	Total Petroleum Hydrocarbons (µg/l)
		Cadmium	Mercury	Lead	Chromium		
SS1	Surface	1.98	<1.0	<1.0	2.24	<0.001	<0.1
	Bottom	2.01	<1.0	<1.0	2.26	<0.001	<0.1
SS2	Surface	1.96	<1.0	<1.0	2.19	<0.001	<0.1
	Bottom	1.99	<1.0	<1.0	2.21	<0.001	<0.1
SS3	Surface	1.95	<1.0	<1.0	2.08	<0.001	<0.1
	Bottom	1.96	<1.0	<1.0	2.10	<0.001	<0.1

Table 5. Seabed Sediment Texture at Bhavnagar creek (November 2021)

Station	D ₅₀ (mm)	Sand (%)				Classification of Sediment
		Coarse Sand	Medium Sand	Fine Sand	Silt & Clay	
SB1	0.11	4.6	9.2	68.6	17.6	Fine Sand
SB2	0.13	4.6	9.5	74.1	11.8	Fine Sand
SB3	0.19	6.8	9.0	69.0	15.2	Fine Sand

Table 6. Seabed Sediment quality parameters at Bhavnagar creek (November 2021)

Station	Total Organic Carbon (%)	Total Nitrogen (mg/kg)	Total Phosphorus (mg/kg)	Calcium Carbonate (%)
SB1	0.68	42.6	16.8	18.4
SB2	0.52	26.8	14.6	21.2
SB3	0.64	39.4	16.4	19.6

Table 7. Concentration of Heavy Metals, Phenols and Total Petroleum Hydrocarbons in Seabed Sediments at Bhavnagar creek (November 2021)

Station	Heavy metals (mg/kg)				Phenols (mg/kg)	Total Petroleum Hydrocarbons (µg/kg)
	Cadmium	Mercury	Lead	Chromium		
SB1	<0.1	<0.1	<0.1	23.2	<0.5	<0.5
SB2	<0.1	<0.1	<0.1	23.8	<0.5	<0.5
SB3	<0.1	<0.1	<0.1	24.6	<0.5	<0.5

Table 8. Primary productivity at Bhavnagar creek water (November 2021)

Station	Gross Photosynthetic activity	Net Photosynthetic activity	Primary production (mgC/m ³ /day)
SS1	1.4	1.1	420
SS2	1.1	0.7	330
SS3	1.2	0.8	360
Average			370

Table 9. Phytoplankton species composition* at Bhavnagar creek water (November 2021)

Sl. No.	Species	Station		
		SS1	SS2	SS3
Class: Bacillariophyceae (Diatoms)				
Order: Centrales				
1	Bellerochea sp.	+	+	+
2	Coscinodiscus Sp.	+	+	+
3	Coscinodiscus grani.	+	+	+
4	Coscinodiscus centralis	+	+	-
5	Hemiaulus sp.	+	-	-
6	Helicotheca thamensis	+	+	+
7	Odontella mobiliensis	-	+	-
8	Odontella sinensis	+	+	+
9	Rhizosolenia sp.	+	+	+
10	Skeletonema sp.	-	+	-
11	Triceratium sp.	+	+	+
Subtotal		9	10	7
Order: Pennales				
12	Bacillaria sp.	+	+	+
13	Bacillaria paradoxa	-	+	-
14	Guinardia striata	+	+	+
15	Navicula sp.	-	+	+
16	Navicula henneydii	+	+	+
17	Nitzschia sp.	+	+	-
18	Nitzschia longissima	-	+	+
19	Pseudo-nitzschia seriata	+	+	+
20	Pleurosigma sp.	-	+	+
21	Pleurosigma directum	+	+	-
22	Gyrosigma sp.	-	+	-
23	Thalassionema nitzschioides	+	-	-
Subtotal		7	11	7
Class: Dinophyceae (Dinoflagellates)				
24	Ceratium Sp.	+	+	+
25	Ceratium fusus	-	+	+

Sl. No.	Species	Station		
		SS1	SS2	SS3
26	<i>Dinophysis</i> sp.	+	-	-
27	<i>Diplopsalopsis</i> sp.	+	+	+
28	<i>Diplopsalopsis meunier</i>	-	+	+
29	<i>Protopteridinium</i> sp.	+	-	-
30	<i>Protopteridinium depressum</i>	-	+	+
Subtotal		4	5	5
Class: Cyanophyceae (Blue-greens)				
31	<i>Pediastrum</i> Sp.	+	+	-
32	<i>Spirogyra</i> Sp.	+	+	+
33	<i>Trichodesmium erythraeum</i>	+	+	+
Subtotal		3	3	2
Grand total		23	29	21

*Net sample

Table 10. Phytoplankton abundance* (cells/l) at Bhavnagar creek water (November 2021)

Sl. No.	Genus / Species	SS1	SS2	SS3	Total	(%)
Phylum: Heterokontophyta						
Class: Bacillariophyceae (Diatoms)						
Order: Centrales 46.8						
1	<i>Belleriochea</i> sp.	50	100	50	200	2.6
2	<i>Coscinodiscus</i> Sp.	100	150	150	400	5.1
3	<i>Coscinodiscus grani</i> .	150	150	0	300	3.8
4	<i>Coscinodiscus centralis</i>	200	250	200	650	8.3
5	<i>Hemiaulus</i> sp.	100	0	0	100	1.3
6	<i>Helicotheca thamensis</i>	200	50	150	400	5.1
7	<i>Odontella mobiliensis</i>	200	50	250	500	6.4
8	<i>Odontella sinensis</i>	100	150	100	350	4.5
9	<i>Rhizosolenia</i> sp.	50	100	100	250	3.2
10	<i>Skeletonema</i> sp.	0	50	0	50	0.6
11	<i>Triceratium</i> sp.	100	250	100	450	5.8
Order: Pennales						
12	<i>Bacillaria</i> sp.	50	100	50	200	2.6
13	<i>Bacillaria paradoxa</i>	150	50	250	450	5.8
14	<i>Guinardia striata</i>	100	150	100	350	4.5

Sl. No.	Genus / Species	SS1	SS2	SS3	Total	(%)
15	<i>Navicula</i> sp.	0	50	50	100	1.3
16	<i>Navicula henneydii</i>	50	100	100	250	3.2
17	<i>Nitzschia</i> sp.	50	50	0	100	1.3
18	<i>Nitzschia longissima</i>	0	100	100	200	2.6
19	<i>Pseudo-nitzschia seriata</i>	100	100	50	250	3.2
20	<i>Pleurosigma</i> sp.	0	50	50	100	1.3
21	<i>Pleurosigma directum</i>	100	50	0	150	1.9
22	<i>Gyrosigma</i> sp.	0	50	0	50	0.6
23	<i>Thalassionema nitzschioides</i>	50	0	0	50	0.6
Percentage of Class: Bacillariophyceae						75.6
Class: Dinophyceae (Dinoflagellates)						
24	<i>Ceratium</i> Sp.	100	200	200	500	6.4
25	<i>Ceratium fusus</i>	0	100	100	200	2.6
26	<i>Dinophysis</i> sp.	50	0	0	50	0.6
27	<i>Diplopsalopsis</i> sp.	50	50	100	200	2.6
28	<i>Diplopsalopsis meunier</i>	0	50	50	100	1.3
29	<i>Protoperidinium</i> sp.	50	0	0	50	0.6
30	<i>Protoperidinium depressum</i>	0	50	50	100	1.3
Percentage of Class: Dinophyceae						15.4
Class: Cyanophyceae (Blue greens)						
31	<i>Pediastrum</i> Sp.	100	50	0	150	1.9
32	<i>Spirogyra</i> Sp.	50	50	100	200	2.6
33	<i>Trichodesmium erythraeum</i>	100	50	200	350	4.5
Percentage of Class - Cyanophyceae						9.0
Grand total		2400	2750	2650	7800	100

* Bottle sample

Table 11. Phytoplankton population at Bhavnagar creek water (November 2021)

Station	No of genera or species (net sample)	Population (cells/l) (bottle sample)
SS1	23	2400
SS2	29	2750
SS3	21	2650

Table 12. Numerical abundance of zooplankton (nos./100m³) at Bhavnagar creek water (November 2021)

Sl. No.	Genus / Species	Station				
		SS1	SS2	SS3	Total	%
Phylum: Protozoa						
Order: Tintinnids (Ciliate groups)						
1	<i>Favella</i> sp.	331	331	0	662	2.3
2	<i>Tintinnopsis</i> sp.	662	993	662	2317	8.0
Phylum: Rotifera						
3	<i>Brachionus plicatilis</i>	662	1325	993	2980	10.3
Phylum: Annelida						
4	Polychaete larvae	331	331	331	993	3.4
Phylum: Arthropoda						
Order: Copepoda						
Sub-order: Calanoida						
5	<i>Acrocalanus</i> sp.	0	331	331	662	2.3
6	<i>Acrocalanus gibber</i>	331	331	0	662	2.3
7	<i>Acartia</i> Sp.	331	0	331	662	2.3
8	<i>Cosmocalanus darwini</i>	0	662	331	993	3.4
9	<i>Clausocalanus thompsoni</i>	662	331	331	1324	4.6
10	<i>Clausocalanus minor</i>	331	331	331	993	3.4
11	<i>Nannocalanus minor</i>	662	993	662	2317	8.0
12	<i>Paracalanus</i> Sp.	331	331	0	662	2.3
13	<i>Temora turbinata</i>	662	331	0	993	3.4
14	Copepod nauplii	1325	993	662	2980	10.3
Sub-order: Cyclopoida						
15	<i>Corycaeus</i> Sp.	331	0	331	662	2.3
16	<i>Oithona</i> sp.	662	331	0	993	3.4
17	<i>Oithona nana</i>	933	1325	993	3251	11.2
Sub-order: Harpacticoida						
18	<i>Microsetella</i> sp.	662	331	0	993	3.4
Other Crustaceans						
19	Shrimp larvae	331	331	662	1324	4.6
20	<i>Lucifer</i> sp.	0	331	331	662	2.3
Phylum: Chordata						
21	Fish larvae	662	331	662	1655	5.7
22	<i>Oikopleura</i> sp.	0	0	331	331	1.1
Total		10202	10594	8275	29071	100

Table 13. Zooplankton diversity, abundance and biomass at Bhavnagar creek water (November 2021)

Station	No. of genera or species	Population (nos./100 m ³)	Biomass (ml/100 m ³)
SS1	18	10202	11.4
SS2	19	10594	13.0
SS3	16	8275	10.7

Table 14. Subtidal and Intertidal benthic population at Bhavnagar creek (November 2021)

Sl. No.	Groups	Subtidal benthic population (nos./m ²)			Intertidal benthic Population (nos./m ²)		
		SB1	SB2	SB3	IB1	IB2	IB3
Phylum: Annelida							
Class: Polychaeta							
1	<i>Armandia sp.</i>	80	40	0	30	0	15
2	<i>Capitella capitella</i>	0	40	0	0	15	0
3	<i>Chaetopterus sp.</i>	40	0	40	15	0	30
4	<i>Dorvillea sp.</i>	80	40	0	0	0	15
5	<i>Cossura sp.</i>	0	40	0	15	0	0
6	<i>Glycera sp</i>	40	0	40	15	30	30
7	<i>Glycinde sp.</i>	0	0	40	0	0	15
8	<i>Perinereis sp.</i>	40	80	0	0	15	0
9	<i>Nereis diversicolor</i>	80	0	40	30	15	0
10	<i>Minuspio cirrifera</i>	40	0	0	15	30	0
11	<i>Prionospio pinnata</i>	0	40	40	0	0	15
12	<i>Pisione indica</i>	0	40	0	15	30	15
Phylum: Arthropoda							
13	Amphipoda	0	40	40	15	15	0
14	Isopoda	40	0	40	0	30	0
Phylum: Mollusca							
15	<i>Cerithidea sp</i>	40	40	0	30	0	15
16	<i>Paphia sp</i>	0	40	40	0	15	15
17	Unidentified bivalves	0	0	40	15	15	0
18	Unidentified gastropods	40	40	0	0	30	15
Total		520	480	360	195	240	180

Table 15. Bacterial population at Bhavnagar creek water ($\times 10^3$ CFU/ml) (November 2021)

Media	Type of Bacteria	Stations		
		SS1	SS2	SS3
Nut Agar	TVC	4.82	5.17	5.18
Mac Agar	TC	0.42	0.39	0.41
Mac Agar	FC	0.31	0.32	0.33
Mac Agar	ECLO	0.16	0.16	0.19
XLD Agar	SHLO	0.17	0.21	0.18
TCBS Agar	VLO	0.16	0.18	0.19
TCBS Agar	VPLO	0.14	0.15	0.17
TCBS Agar	VCLO	0.01	0.1	0.04

Table 16. Bacterial population at Bhavnagar creek sediments ($\times 10^4$ CFU/g) (November 2021)

Media	Type of Bacteria	Stations		
		SB1	SB2	SB3
Nut Agar	TVC	5.15	5.2	5.23
Mac Agar	TC	49	0.52	0.57
Mac Agar	FC	0.31	0.3	0.34
Mac Agar	ECLO	0.15	0.18	0.17
XLD Agar	SHLO	0.19	0.2	0.24
TCBS Agar	VLO	0.19	0.18	0.19
TCBS Agar	VPLO	0.15	0.16	0.18
TCBS Agar	VCLO	0.02	0.02	0.03

TVC - Total Viable Counts; TC - Total Coliforms; FC- Faecal coliform; ECLO - *Escherichia coli* like organisms; SHLO - *Shigella* like organisms; VLO - *Vibrio* like organisms; VPLO - *Vibrio parahaemolyticus* like organisms; VCLO-*Vibrio cholerae* like organisms.

Table17. Comparison of Water quality parameters between Pre-Project period (May 2013) and Monitoring period (November 2021)

Sl. No.	Parameters	Unit	Pre-project period	Monitoring period				Remarks
			May 2013	February 2021	May 2021	August 2021	November 2021	
			Range	Range	Range	Range	Range	
1	Temperature	(°C)	32-34	26.5-26.8	Due to COVID-19 Nationwide lockdown and strict interstate travel protocols, quarterly monitoring study for the 1 st season period March 2021 – May 2021 was not conducted	27.0-27.6	26.0-27.0	Seasonal variation
2	pH	-	8.0-8.2	7.88-7.97		7.23-7.48	7.34-7.49	Seasonal variation
3	Salinity	ppt	36-37	32.5-33.4		27.5-29.5	19.1-20.1	Seasonal variation
4	Turbidity	NTU	>1000	610-880		570-661	691-812	Seasonal variation
5	TSS	mg/l	980-13052	1270-1584		1254-1454	1520-1786	Seasonal variation
6	DO	mg/l	3.84-4.32	5.35-5.6		5.2-5.8	5.8-6.1	No significant change
7	BOD	mg/l	1.44-3.20	0.9-1.3		1.0-1.3	1.2-1.6	No significant change
8	COD	mg/l	37.3-52.5	18.8-21.6		20.4-24	25.6-33.6	Seasonal variation
9	Ammoniacal Nitrogen	µmol/l	0.43-0.74	1.5-2.2		2.61-3.27	1.61-2.27	Seasonal variation
10	Nitrite	µmol/l	1.56-3.47	0.53-0.91		1.54-2.11	1.20-1.34	Seasonal variation
11	Nitrate	µmol/l	3.76-7.78	3.12-4.10		14.2-16.6	4.2-6.6	Seasonal variation
12	Total Nitrogen	µmol/l	13.92-26.36	5.55-7.25		18.8-21.4	7.55-9.94	Seasonal variation
13	Phosphate	µmol/l	0.38-2.77	0.79-0.97		1.42-1.74	1.5-2.3	Seasonal variation
14	Total Phosphorus	µmol/l	5.23-7.22	2.53-3.35		2.23-3.1	2.4-3.4	Seasonal variation
15	Cadmium	µg/l	<1.0	1.8-2.2		2.1-2.6	1.95-2.01	No significant change
16	Mercury	µg/l	<1.0	<1.0		<1.0	<1.0	No change
17	Lead	µg/l	<1.0	<1.0		<1.0	<1.0	No change
18	Chromium	µg/l	2.95-5.64	2.24-2.48		2.1-2.5	2.08-2.26	Seasonal variation
19	Phenols	mg/l	<0.001	<0.001		<0.001	<0.001	No change
20	Total Petroleum Hydrocarbons	µg/l	<0.05	<0.1		<0.1	<0.1	No change

*For May 2021, samples was not collected due to nationwide ban on travel due to COVID-19 pandemic.

Table 18. Comparison of Sediment quality parameters between Pre-Project period (May 2013) and Monitoring period (November 2021)

Sl. No.	Parameters	Unit	Pre-project period	Monitoring period				Remarks
			May 2013	February 2021	May 2021	August 2021	November 2021	
			Range	Range	Range	Range	Range	
1	Soil texture	-	Fine sand	Fine Sand	Due to COVID-19 Nationwide lockdown and strict interstate travel protocols, quarterly monitoring study for the 1 st season period March 2021 – May 2021 was not conducted	Fine Sand	Fine Sand	Seasonal variation
2	Total Phosphorous	mg/kg	0.26-0.38	15.8-18.6		12.6-15.2	14.6-16.8	Seasonal variation but within normal range
3	Total Nitrogen	mg/kg	0.86-1.22	16.4-26.4		21.2-28.6	26.8-42.6	Seasonal variation but within normal range
4	Total Organic Carbon	%	0.47-0.77	0.20-1.17		0.46-0.62	0.52-0.68	Seasonal variation but within normal range
5	Calcium Carbonate	%	5.76-7.72	12.5-20.6		16.8-24.4	18.4-21.2	Seasonal variation but within normal range
6	Cadmium	mg/kg	<1.0	<0.1		<0.1	<0.1	No change
7	Chromium	mg/kg	30.48-35.08	22.4-32.6		23.8-25.4	23.2-24.6	Marginal change but within normal range
8	Mercury	mg/kg	<1.0	<0.1		<0.1	<0.1	No change
9	Lead	mg/kg	<1.0	<0.1		<0.1	<0.1	No change
10	Phenols	mg/kg	<0.05	<0.5		<0.5	<0.5	No change
11	Petroleum Hydrocarbons	µg/kg	2.12-3.37	<0.5		<0.5	<0.5	No change

Table 19. Comparison of Biological parameters between Pre-Project period (May 2013) and Monitoring period (November 2021)

Sl. No.	Parameters	Unit	Pre-project period	Monitoring period				Remarks
			May 2013	February 2021	May 2021	August 2021	November 2021	
Phytoplankton								
1	Primary Productivity	mgC/m³/day	240-480	330-420	Due to COVID-19 Nationwide lockdown, 1 st season period March 2021 – May 2021 was not conducted	330-390	330-420	No significant change
2	Species composition	nos./100m³	12-31	16-21		21-25	21-29	Seasonal variation
3	Abundance	nos./l	399-2530	1450-2900		1950-2750	2400-2750	No significant change
Zooplankton								
4	Biomass	ml/100m³	5.14-14.85	9.6-11.7		10.6-12.2	10.7-13.0	No significant change
5	Abundance	nos./100m³	7368-50048	7159-10034		8300-11840	8275-10594	Seasonal variation
Benthos								
6	Subtidal	nos./m²	20-30	280-400		360-480	360-520	No significant change
7	Intertidal	nos./m²	20-40	150-210		165-240	180-240	No significant change
Microbial count								
8	Water	nos. x10³/ml	0.01-5.31	0.02-5.02		0.01-5.21	0.01-5.18	No significant change
9	Sediment	nos. x10⁴/g	0.01-5.48	0.01-5.21		0.02-5.24	0.02-5.23	No significant change



Collection of water sample



Onboard testing



Collection of sediment sample



Collection of plankton sample



Collection of Intertidal benthos sample

Annexure I – QCI NABET accreditation certificate



Quality Council of India
**National Accreditation Board for
Education & Training**

CERTIFICATE OF ACCREDITATION

Indomer Coastal Hydraulics (P) Ltd.

63, Gandhi road, Alwarthirunagar, Chennai 600087

The organization is accredited as Category-A under the QCI-NABET Scheme for Accreditation of EIA Consultant Organization, Version 3: for preparing EIA-EMP reports in the following Sectors –

Sl. No.	Sector Description	Sector (as per)		Cat.
		NABET	MoEFCC	
1.	Oil & gas transportation pipeline (crude and refinery/ petrochemical products), passing through national parks/ sanctuaries/coral reefs / ecologically sensitive areas including LNG terminal	27	6 (a)	A
2.	Ports, harbours, break waters and dredging	33	7 (e)	A

Note: Names of approved EIA Coordinators and Functional Area Experts are mentioned in RAAC and Supplementary MoM dated Jan 27, 2021, and June 08, 2021 respectively posted on QCI-NABET website.

The Accreditation shall remain in force subject to continued compliance to the terms and conditions mentioned in QCI-NABET's letter of accreditation bearing no. QCI/NABET/ENV/ACO/21/1784 dated June 29, 2021. The accreditation needs to be renewed before the expiry Indomer Coastal Hydraulics (P) Ltd. following due process of assessment.



Sr. Director, NABET
Dated: June 29, 2021

Certificate No.
NABET/EIA/2023/RA 0207

Valid till
Sept 13, 2023

For the updated List of Accredited EIA Consultant Organizations with approved Sectors please refer to QCI-NABET website.

Annexure II – NABL accreditation certificate



National Accreditation Board for
Testing and Calibration Laboratories

CERTIFICATE OF ACCREDITATION

INDOMER COASTAL HYDRAULICS PRIVATE LIMITED

has been assessed and accredited in accordance with the standard

ISO/IEC 17025:2017

**"General Requirements for the Competence of Testing &
Calibration Laboratories"**

for its facilities at

NO 63, GANDHI ROAD, ALWAR THIRUNAGAR, CHENNAI, TAMIL NADU, INDIA

in the field of

TESTING

Certificate Number: TC-5232

Issue Date: 13/01/2019

Valid Until: 12/01/2021*

* The validity is extended for one year up to 12.01.2022

*Transition to 2017 version completed w.e.f 22.04.2021 valid until 12.01.2022

This certificate remains valid for the Scope of Accreditation as specified in the annexure subject to continued satisfactory compliance to the above standard & the relevant requirements of NABL.
(To see the scope of accreditation of this laboratory, you may also visit NABL website www.nabl-india.org)

Name of Legal Identity : INDOMER COASTAL HYDRAULICS PRIVATE LIMITED

Signed for and on behalf of NABL



N. Venkateswaran
Chief Executive Officer




FIG.1. LOCATION MAP




FIG. 2 . SAMPLING LOCATIONS